

A GKM DESCRIPTION OF THE EQUIVARIANT COHOMOLOGY RING OF A HOMOGENEOUS SPACE

V. GUILLEMIN, T. HOLM, AND C. ZARA

ABSTRACT. Let T be a torus of dimension $n > 1$ and M a compact T -manifold. M is a GKM manifold if the set of zero dimensional orbits in the orbit space M/T is zero dimensional and the set of one dimensional orbits in M/T is one dimensional. For such a manifold these sets of orbits have the structure of a labelled graph and it is known that a lot of topological information about M is encoded in this graph.

In this paper we prove that every compact homogeneous space M of non-zero Euler characteristic is of GKM type and show that the graph associated with M encodes *geometric* information about M as well as topological information. For example, from this graph one can detect whether M admits an invariant complex structure or an invariant almost complex structure.

CONTENTS

1. Introduction	2
2. The Equivariant Cohomology of Homogeneous Spaces	4
2.1. The Borel description	4
2.2. The GKM graph of M	4
2.3. The GKM definition of the cohomology ring	9
2.4. Equivalence between the Borel picture and the GKM picture	10
3. Almost complex structures and axial functions	12
3.1. Axial functions	12
3.2. Invariant almost complex structures	13
4. Morse theory on the GKM graph	14
4.1. Betti numbers	14
4.2. Morse functions	15
4.3. Invariant complex structures	15
5. Examples	16
5.1. Non-existence of almost complex structures	16
5.2. Non-existence of Morse functions	17
5.3. The existence of several almost complex structures	17
References	18

1. INTRODUCTION

Let T be a torus of dimension $n > 1$, M a compact manifold,

$$\tau : T \times M \rightarrow M$$

a faithful action of T on M , and M/T the orbit space of τ . M is called a *GKM manifold* if the set of zero dimensional orbits in the orbit space M/T is zero dimensional and the set of one dimensional orbits in M/T is one dimensional. Under these hypotheses, the union, $\Gamma \subset M/T$, of the set of zero and one dimensional orbits has the structure of a graph: Each connected component of the set of one-dimensional orbits has at most two zero-dimensional orbits in its closure; so these components can be taken to be the edges of a graph and the zero-dimensional orbits to be the vertices. Moreover, each edge, e , of Γ consists of orbits of the same orbittype: namely, orbits of the form $\mathcal{O}_e = T/H_e$, where H_e is a codimension one subgroup of T . Hence one has a labelling

$$e \rightarrow H_e \tag{1.1}$$

of the edges of Γ by codimension one subgroups of T .

It has recently been discovered that if M has either a T -invariant complex structure or a T -invariant symplectic structure, the data above - the graph Γ and the labelling (1.1) - contain a surprisingly large amount of information about the global topology of M . For instance, Goresky, Kottwitz, and MacPherson proved that the ring structure of the equivariant cohomology ring $H_T^*(M)$ is completely determined by this data, and Knutson and Rosu have shown that the same is true for the ring $K_T(M) \otimes \mathbb{C}$.

The manifolds M which we will be considering below will be neither complex nor symplectic; however we will make an assumption about them which is in some sense much stronger than either of these assumptions. We will assume that T is the Cartan subgroup of a compact, semisimple, connected Lie group G , and that G acts transitively on M , *i.e.* M is a G -homogeneous space. There is a simple criterion for such a manifold to be a GKM manifold.

Theorem 1.1. *Suppose M is a G -homogeneous manifold. Then the following are equivalent.*

1. *The action of T on M is a GKM action;*
2. *The Euler characteristic of M is non-zero;*
3. *M is of the form $M = G/K$, where K is a closed subgroup of G containing T .*

As we mentioned above, the data (1.1) determine the ring structure of $H_T^*(M)$ if M is either complex or symplectic. This result is, in fact, true modulo an assumption which is weaker than either of these assumptions; and this assumption - equivariant formality - is satisfied by homogeneous spaces which satisfy the hypotheses of the theorem. Hence, for these spaces, one has two completely different descriptions of the ring $H_T^*(M)$: the graph

theoretical description above and the classical Borel description. In Section 2 we will compute the graph Γ of a space M of the form G/K , with $T \subset K$, and show that it is a *homogeneous* graph, i.e. we will show that the Weyl group of G , W_G , acts transitively on the vertices of Γ and that this action preserves the labelling (1.1). We will then use this result to compare the two descriptions of $H_T^*(M)$.

One of the main goals in this paper is to show that for homogeneous manifolds M of GKM type, some important features of the geometry of M can be discerned from the graph Γ and the labelling (1.1). One such feature is the existence of a G -invariant almost complex structure. The subgroups, H_e , labelling the edges of Γ are of codimension one in T ; so, up to sign, they correspond to weights, α_e , of the group T . It is known that the W_K -invariant labelling (1.1) can be lifted to a W_K -invariant labelling

$$e \rightarrow \alpha_e \tag{1.2}$$

if M is a coadjoint orbit of G (hence, in particular, a complex G -manifold). Moreover, this labelling has certain simple properties which we axiomatize by calling a map with these properties an *axial function* (see Section 3.1). In Section 3 we prove the following result.

Theorem 1.2. *The homogeneous space M admits a G -invariant almost complex structure if and only if Γ possesses a W_K -invariant axial function (1.2) compatible with (1.1).*

This raises the issue: Is it possible to detect from the graph theoretic properties of the axial function (1.2) whether or not M admits a G -invariant complex structure? Fix a vector $\xi \in \mathfrak{t}$ such that $\alpha_e(\xi) \neq 0$ for all oriented edges, e , of Γ , and orient these edges by requiring that $\alpha_e(\xi) > 0$. We prove in Section 4 the following theorem.

Theorem 1.3. *A necessary and sufficient condition for M to admit a G -invariant complex structure is that there exist no oriented cycles in Γ .*

Remarks:

1. M admits a G -invariant complex structure if and only if it admits a G -invariant symplectic structure; and, by the Konstant-Kirillov theorem, it has either (and hence both) of these properties if and only if it is a coadjoint orbit of G .
2. By the Goresky-Kottwitz-MacPherson theorem, the graph Γ and the axial function (1.2) determine the cohomology ring structure of M . The *additive* cohomology of M , i.e. its Betti numbers, β_i , can be computed by the following simple recipe: For each vertex, p , of the graph Γ , let σ_p be the number of oriented edges issuing from p with the property that $\alpha_e(\xi) < 0$. Then

$$\beta_i = \begin{cases} 0, & \text{if } i \text{ is odd,} \\ \#\{p; \sigma_p = i/2\}, & \text{if } i \text{ is even.} \end{cases}$$

One question we have not addressed in this paper is the question: When is a labelled graph the GKM graph of a homogeneous space of the form G/K with $T \subset K$? For some partial answers to this question see [Ho].

Acknowledgements: We are grateful to David Vogan for helping us formulate and prove the results in Section 3 and to Bert Kostant for pointing out to us that a homogeneous space is a quotient of a compact group by a closed subgroup of the same rank if and only if its Euler characteristic is non-zero, and for making us aware of a number of nice properties of such spaces.

2. THE EQUIVARIANT COHOMOLOGY OF HOMOGENEOUS SPACES

2.1. The Borel description. Let G be a compact semi-simple Lie group, T a Cartan subgroup of G , K a closed subgroup of G such that

$$T \subset K \subset G ,$$

and let $\mathfrak{t} \subset \mathfrak{k} \subset \mathfrak{g}$ be the Lie algebras of T , K , and G .

Let $\Delta_K \subset \Delta_G$ be the roots of K and G , with $\Delta_K^+ \subset \Delta_G^+$ sets of positive roots, let

$$\Delta_{G,K} = \Delta_G - \Delta_K ,$$

and let $W_K \subset W_G$ be the Weyl groups of K and G . We will regard an element of W_G both as an element of $N(T)/T$ and as a transformation of the dual Lie algebra $\mathfrak{t}^*$ (or as a transformation of $\mathfrak{t}$, via the isomorphism $\mathfrak{t}^* \simeq \mathfrak{t}$ given by the Killing form). Also, we will assume for simplicity that G is simply connected and that the homogeneous space G/K is oriented.

Now suppose M is a G -manifold. Then the equivariant cohomology ring $H_T^*(M)$ is related to the cohomology ring $H_G^*(M)$ by

$$H_T^*(M) = H_G^*(M) \otimes_{\mathbb{S}(\mathfrak{t}^*)^{W_G}} \mathbb{S}(\mathfrak{t}^*) .$$

(see [GS, Chap. 6]), where $\mathbb{S}(\mathfrak{t}^*)$ is the symmetric algebra of $\mathfrak{t}^*$. In particular, let $M = G/K$, where K acts on G by *right* multiplication. Then G acts on M by *left* multiplication and

$$H_G^*(M) = H_G^*(G/K) = \mathbb{S}(\mathfrak{t}^*)^K = \mathbb{S}(\mathfrak{t}^*)^{W_K} ,$$

hence

$$H_T^*(G/K) = \mathbb{S}(\mathfrak{t}^*)^{W_K} \otimes_{\mathbb{S}(\mathfrak{t}^*)^{W_G}} \mathbb{S}(\mathfrak{t}^*) . \quad (2.1)$$

This is the Borel description of $H_T^*(G/K)$. Throughout this paper, unless stated otherwise, M is the homogeneous space G/K .

2.2. The GKM graph of M . In the following subsections, we will show that M is equivariantly formal and is a GKM space. Then we will relate the the GKM description of the equivariant cohomology ring of M to the description above.

2.2.1. *Equivariant formality.* The $\mathbb{S}(\mathfrak{t}^*)$ -module structure of the equivariant cohomology ring $H_T^*(M)$ can be computed by a spectral sequence (see [GS, p. 70]) whose E_2 term is $H(M) \otimes \mathbb{S}(\mathfrak{t}^*)$, and if this spectral sequence collapses *at this stage*, then M is said to be *equivariantly formal*. If $M = G/K$, with $T \subset K$, then,

$$H^{odd}(M) = 0,$$

(see [GHV3, p. 467]), and from this it is easy to see that all the higher order coboundary operators in this spectral sequence have to vanish by simple degree considerations. Hence M is equivariantly formal. One implication of equivariant formality is the following:

Theorem 2.1. *The restriction map*

$$H_T^*(M) \rightarrow H_T^*(M^T) \tag{2.2}$$

is injective.

Proof. By a localization theorem of Borel (see [Bo]), the kernel of (2.2) is the torsion submodule of $H_T^*(M)$. However, if M is equivariantly formal, then $H_T^*(M)$ is free as an $\mathbb{S}(\mathfrak{t}^*)$ -module, so the kernel has to be zero. $\square$

Thus $H_T^*(M)$ imbeds as a subring of the ring

$$H_T^*(M^T) = H^0(M^T) \otimes_{\mathbb{C}} \mathbb{S}(\mathfrak{t}^*) . \tag{2.3}$$

We will give an explicit description of this subring in Section 2.3.

2.2.2. *The Euler characteristic.* It follows from (2.3) that, if M is a homogeneous space of the form G/K , with $T \subseteq K$, then the Euler characteristic of M is equal to

$$\chi(M) = \sum_i \dim H^{2i}(M) ;$$

in particular, the Euler characteristic is non-zero. It is easy to see that the converse is true as well.

Proposition 2.1. *If $M = G/K$ and the rank of K is strictly less than the rank of G , then the Euler characteristic of G/K is zero.*

Proof. Let h be an element of T with the property that

$$\{h^N ; -\infty < N < \infty\}$$

is dense in T . Suppose that the action of h on G/K fixes a coset g_0K . Then $g_0^{-1}hg_0 \in K$, i.e. h is conjugate to an element of K and hence conjugate to an element h_1 of the Cartan subgroup T_1 of K . However, if the iterates of h are dense in T , so must be the iterates of h_1 and hence $T_1 = T$. Suppose now that $h = \exp \xi, \xi \in \mathfrak{t}$. If h has no fixed points, then the vector field ξ_M can have no zeroes and hence the Euler characteristic of M has to be zero. $\square$

2.2.3. The fixed points. We prove in this section that the action of T on M is a GKM action; i.e. that the set of zero dimensional orbits in the orbit space M/T is zero dimensional, and the set of one dimensional orbits is one dimensional. It is easy to see that these properties are equivalent to

- (1) M^T is finite;
- (2) For every codimension one subgroup H of T , $\dim M^H \leq 2$.

We will show that if M is of the form G/K , with $T \subseteq K$, then it has the two properties above, and we will also show that it has the following third property:

- (3) For every subtorus H of T and every connected component X of M^H , $X^T \neq \emptyset$.

It is well known that these properties hold for the homogeneous space $\mathcal{O} = G/T$. The first two properties can be checked directly (see [GZ1], and the third property holds because $\mathcal{O}$ is a compact symplectic manifold and the action of T is Hamiltonian. Therefore, to prove that M satisfies properties 1-3, it suffices to prove the following theorem.

Theorem 2.2. *For every subtorus H of T , the map*

$$\mathcal{O} = G/T \rightarrow G/K = M \tag{2.4}$$

sends $\mathcal{O}^H$ onto M^H .

Proof. Let p_0 be the identity coset in M and q_0 the identity coset in $\mathcal{O}$. Let h be an element of H with the property that

$$\{h^N ; -\infty < N < \infty\}$$

is dense in H . If $p = gp_0 \in M^H$, then $g^{-1}hg \in K$; so $g^{-1}hg = ata^{-1}$, with $a \in K$ and $t \in T$. Thus $hga = gat$ and hence $hq = q$, where $q = gaq_0$. But under the map (2.4), q_0 is sent to p_0 , so q is sent to $gap_0 = gp_0 = p$. $\square$

In particular, Theorem 2.2 tells us that the map $\mathcal{O}^T \rightarrow M^T$ is surjective. However,

$$\mathcal{O}^T = N_G(T)/T = W_G ,$$

so M^T is the image of $W_G = N_G(T)/T$ in G/K . But $N_G(T) \cap K = N_K(T)$, the normalizer of T in K , so

$$(N_G(T) \cap K)/T = W_K ,$$

and hence we proved:

Proposition 2.2. *There is a bijection*

$$M^T \simeq W_G/W_K ;$$

in particular, $W_G = N_G(T)/T$ acts transitively on M^T .

2.2.4. Points stabilized by codimension one subgroups. Next we compute the connected components of the sets M^H , where H is a codimension one subgroup of T . Let X be one of these components. Then $X^T \neq \emptyset$. Moreover, since M is simply connected, it is orientable, and hence every connected component of M^H is orientable. So, if X is not an isolated point of M^H , then it has to be either a circle, a 2-torus, or a 2-sphere, and the first two possibilities are ruled out by the condition $X^T \neq \emptyset$. We conclude:

Theorem 2.3. *Let H be a codimension one subgroup of T and let X be a connected component of M^H . Then X is either a point or a 2-sphere.*

Remark. By the Korn-Lichtenstein theorem, every faithful action of S^1 on the 2-sphere is diffeomorphic to the standard action of “rotation about the z -axis”. Therefore the action of the circle $S^1 = T/H$ on the 2-sphere X in the theorem above has to be diffeomorphic to the standard action. In particular, $\#X^T = 2$.

We now explicitly determine what these 2-spheres are. By Theorem 2.2, each of these 2-spheres is the conjugate by an element of $N_G(T)$ of a 2-sphere containing the identity coset $p_0 \in M = G/K$; so we begin by determining the 2-spheres containing p_0 .

2.2.5. The space $\mathfrak{g}/\mathfrak{k}$. The tangent space $T_{p_0}M$ can be identified with $\mathfrak{g}/\mathfrak{k}$, and the isotropy representation of T on this space decomposes into a direct sum of two-dimensional T -invariant subspaces

$$T_{p_0}M = \bigoplus V_{[\alpha]} , \quad (2.5)$$

labelled by the roots modulo ± 1 ,

$$\alpha \in \Delta_{G,K} / \pm 1 . \quad (2.6)$$

One can also regard this as a labelling by the *positive* roots in $\Delta_{G,K}$; however, since this set of positive roots is not fixed by the natural action of W_K on $\Delta_{G,K}$, this is not an *intrinsic* labelling. (This fact is of importance in Section 3, when we discuss the existence of G -invariant almost complex structures on M .) Now let H be a codimension one subgroup of T , let $\mathfrak{h} \subset \mathfrak{t}$ be the Lie algebra of H , and let M^H be the set of H -fixed points. Then

$$T_{p_0}M^H = (T_{p_0}M)^H .$$

Hence, if X is the connected component of M^H containing p_0 , and if X is not an isolated point, then $(T_{p_0}M)^H$ has to be one of the $V_{[\alpha]}$'s in the sum (2.5). Hence the adjoint action of H on $\mathfrak{g}/\mathfrak{k}$ has to leave $V_{[\alpha]}$ pointwise fixed. However, an element $g = \exp t$ of T acts on $V_{[\alpha]}$ by the rotation

$$\chi_\alpha(g) = \begin{pmatrix} \cos \alpha(t) & -\sin \alpha(t) \\ \sin \alpha(t) & \cos \alpha(t) \end{pmatrix} , \quad (2.7)$$

so the stabilizer group of $V_{[\alpha]}$ is the group

$$H_\alpha = \{g \in T ; \chi_\alpha(g) = 1\} . \quad (2.8)$$

Let $C(H_\alpha)$ be the centralizer of H_α in G and let G_α be the semisimple component of $C(H_\alpha)$. Then G_α is either $SU(2)$ or $SO(3)$, and since G_α is contained in $C(H_\alpha)$, $G_\alpha p_0$ is fixed pointwise by the action of H . Moreover, since $G_\alpha \not\subseteq K$, the orbit $G_\alpha p_0$ can't just consist of the point p_0 itself; hence

$$G_\alpha p_0 = X. \quad (2.9)$$

The Weyl group of G_α is contained in the Weyl group of G and consists of two elements: the identity and a reflection, $\sigma = \sigma_\alpha$, which leaves fixed the hyperplane $\ker \alpha \subset \mathfrak{t}$, and maps α to $-\alpha$. Therefore, since $\alpha \notin \Delta_K$, $\sigma_\alpha p_0 \neq p_0$, and hence p_0 and $\sigma_\alpha p_0$ are the two T -fixed points on the 2-sphere (2.9).

Now let $p = wp_0$ be another fixed point of T , with $[w] \in W_G/W_K$. Let a be a representative for w in $N_G(T)$ and let $L_a : G \rightarrow G$ be the left action of a on G . If X is the 2-sphere (2.9), then the 2-sphere $L_a(X)$ intersects M^T in the two fixed points wp_0 and $w\sigma_\alpha p_0$, and its stabilizer group in T is the group

$$aH_\alpha a^{-1} = wH_\alpha w^{-1} = H_{w\alpha}, \quad (2.10)$$

where H_α is the group (2.8).

2.2.6. The GKM graph of M . This concludes our classification of the set of 2-spheres in M which are stabilized by codimension one subgroups of T . Now note that if X is such a two-sphere and H is the subgroup of T stabilizing it, then the orbit space X/T consists of two T -fixed points and a connected one dimensional set of orbits having the orbittype of T/H . Thus these X 's are in one-to-one correspondence with the edges of the GKM graph of M . Denoting this graph by Γ we summarize the graph-theoretical content of what we've proved so far:

Theorem 2.4. *The GKM data associated to the action of T on the homogeneous space $M = G/K$ is the following.*

- (1) *The vertices of Γ are in one-to-one correspondence with the elements of W_G/W_K ;*
- (2) *Two vertices $[w]$ and $[w']$ are on a common edge of Γ if and only if $[w'] = [w\sigma_\alpha]$ for some $\alpha \in \Delta_{G,K}$;*
- (3) *The edges of Γ containing the vertex $[w]$ are in one-to-one correspondence with the roots, modulo ± 1 , in the set $\Delta_{G,K}$;*
- (4) *If α is such a root, then the stabilizer group (1.1) labelling the edge corresponding to this root is the group (2.10).*

In particular, the labelling (1.1) of the graph Γ can be viewed as a labelling by elements $[\alpha]$ of $\Delta_G/\pm 1$. We call this labelling a *pre-axial function*.

2.2.7. The connection on Γ . One last structural component of the graph Γ remains to be described: Given any graph, Γ , and vertex, p , of Γ , let E_p be

the set of *oriented* edges of Γ with initial vertex p . A *connection* on Γ is a function which assigns to each oriented edge, e , a bijective map

$$\theta_e : E_p \rightarrow E_q ,$$

where p is the initial vertex of e and q is the terminal vertex. The graph Γ described in Theorem 2.4 has a natural such connection. Namely, let e be the oriented edge of Γ joining $[w]$ to $[w\sigma_\alpha]$. If $e' \in E_{[w]}$ is the oriented edge joining $[w]$ to $[w\sigma_\beta]$, then let $\theta_e(e') = e''$, where e'' is the edge joining $[w\sigma_\alpha]$ to $[w\sigma_\alpha\sigma_\beta]$. This connection is compatible with the pre-axial function (1.1) in the sense that, for every vertex p , and every pair of oriented edges, $e, e' \in E_p$, the roots labelling e, e' , and $e'' = \theta_e(e')$ are coplanar in $\mathfrak{t}^*$.

2.2.8. Simplicity. A graph is said to be *simple* if every pair of vertices is joined by at most one edge. Most of the graphs above don't have this property. There is however an important class of subgroups, K , for which the graph associated with G/K does have this property.

Theorem 2.5. *If K is the stabilizer group of an element of $\mathfrak{t}$, then the graph Γ is simple.*

Proof. A root $\alpha \in \Delta_G$ is in Δ_K if and only if the restriction of α to the subspace $\mathfrak{t}^{W_K}$ of $\mathfrak{t}$ is zero. Let $\alpha, \beta \in \Delta_{G,K}$ such that $\alpha \neq \pm\beta$, and let $\sigma_\alpha, \sigma_\beta$ be the reflections of $\mathfrak{t}$ defined by α and β . Then $\sigma_\alpha \neq \sigma_\beta$ and the subspace of $\mathfrak{t}$ fixed by $\sigma_\alpha\sigma_\beta$ is the codimension 2 subspace on which both α and β vanish. If $\sigma_\alpha\sigma_\beta \in W_K$, then this subspace contains $\mathfrak{t}^{W_K}$, so α and β are both vanishing on $\mathfrak{t}^{W_K}$, contradicting our assumption that $\alpha, \beta \notin \Delta_K$. $\square$

Another way to prove Theorem 2.5 is to observe that $M = G/K$ is a coadjoint orbit of the group G . In particular, it is a Hamiltonian T -space and Γ is the one-skeleton of its moment polytope.

2.3. The GKM definition of the cohomology ring. We recall how the data encoded in the GKM graph determines the equivariant cohomology ring $H_T^*(M)$. The inclusion $i : M^T \rightarrow M$ induces a map in cohomology

$$i^* : H_T^*(M) \rightarrow H_T^*(M^T) = \text{Maps}(M^T, \mathbb{S}(\mathfrak{t}^*)) = \text{Maps}(W_G/W_K, \mathbb{S}(\mathfrak{t}^*)) ,$$

and the fact that M is equivariantly formal implies that i^* is injective. Let $H_T^*(\Gamma)$ be the set of maps

$$f : W_G/W_K \rightarrow \mathbb{S}(\mathfrak{t}^*) \tag{2.11}$$

that satisfy the compatibility condition:

$$f([w\sigma_\alpha]) - f([w]) \in (w\alpha)\mathbb{S}(\mathfrak{t}^*) . \tag{2.12}$$

for every edge $([w], [w\sigma_\alpha])$ of Γ .

The Goresky, Kottwitz and MacPherson theorem [GKM] asserts that

$$H_T^*(M) \simeq i^*(H_T^*(M)) = H_T^*(\Gamma) .$$

In the next section we construct a direct isomorphism between this ring $H_T^*(M)$ and the Borel ring given in (2.1).

2.4. Equivalence between the Borel picture and the GKM picture.

From the inclusion, i , of M^T into M , one gets a restriction map

$$i^* : H_T^*(M) \rightarrow H_T^*(M^T) ; \quad (2.13)$$

and, since M is equivariantly formal, i^* maps $H_T^*(M)$ bijectively onto the subring $H_T^*\Gamma$ of $H_T^*(M^T)$. However, as we pointed out in Section 2.1,

$$H_T^*(M) \simeq \mathbb{S}(\mathfrak{t}^*)^{W_K} \otimes_{\mathbb{S}(\mathfrak{t}^*)^{W_G}} \mathbb{S}(\mathfrak{t}^*) ;$$

so, by combining (2.13) and (2.1), we get an isomorphism

$$\mathcal{K} : \mathbb{S}(\mathfrak{t}^*)^{W_K} \otimes_{\mathbb{S}(\mathfrak{t}^*)^{W_G}} \mathbb{S}(\mathfrak{t}^*) \rightarrow H_T^*(\Gamma) . \quad (2.14)$$

The purpose of this section is to give an explicit formula for this map. Note that since M^T is a finite set,

$$H_T^*(M^T) = \bigoplus_{p \in M^T} H_T^*(p) = \bigoplus_{p \in M^T} \mathbb{S}(\mathfrak{t}^*) = \text{Maps}(M^T, \mathbb{S}(\mathfrak{t}^*)) .$$

Theorem 2.6. *On decomposable elements, $f_1 \otimes f_2$, of the product (2.1),*

$$\mathcal{K}(f_1 \otimes f_2) = g \in \text{Maps}(M^T, \mathbb{S}(\mathfrak{t}^*)) , \quad (2.15)$$

where, for $w \in W_G$ and $p = wp_0 \in M^T$,

$$g(wp_0) = (wf_1)f_2 . \quad (2.16)$$

Proof. We first show that (2.15) and (2.16) do define a ring homomorphism of the ring (2.1) into $H^*(\Gamma)$. To show that (2.16) doesn't depend on the representative w chosen, we note that if $wp_0 = w'p_0$, then $\sigma = w(w')^{-1} \in W_K$. Thus

$$g(w'p_0) = (w'f_1)f_2 = (w\sigma f_1)f_2 = (wf_1)f_2 = g(wp_0) ,$$

since $f_1 \in \mathbb{S}(\mathfrak{t}^*)^{W_K}$. Next, we note that if $f \in \mathbb{S}(\mathfrak{t}^*)^{W_G}$, then

$$\mathcal{K}(f_1 f \otimes f_2) = \mathcal{K}(f_1 \otimes f f_2) ,$$

since

$$w(f_1 f)f_2 = (wf_1)(wf)f_2 = (wf_1)f f_2 .$$

Thus, by the universality property of tensor products, $\mathcal{K}$ does extend to a mapping of the ring (2.1) into the ring $\text{Maps}(M^T, \mathbb{S}(\mathfrak{t}^*))$. Next, let α be a root and let $\sigma \in W_G$ be the reflection that interchanges α and $-\alpha$ and that is the identity on the hyperplane

$$\mathfrak{h} = \{\xi \in \mathfrak{t} ; \alpha(\xi) = 0\} .$$

Suppose that p and p' are two adjacent vertices of Γ with $p' = \sigma p$. To show that $g = \mathcal{K}(f_1 \otimes f_2)$ is in $H_T^*(\Gamma)$, we must show that the quotient

$$\frac{g(p') - g(p)}{\alpha}$$

is in $\mathbb{S}(\mathfrak{t}^*)$. However, if $p = wp_0$, then

$$g(p') - g(p) = (\sigma w f_1 - w f_1)f_2 ,$$

and since σ is the identity on $\mathfrak{h}$, the restriction of the polynomial $w\sigma f_1$ to $\mathfrak{h}$ is equal to the restriction of the polynomial wf_1 to $\mathfrak{h}$; hence

$$\frac{g(p') - g(p)}{\alpha} \in \mathbb{S}(\mathfrak{t}^*) .$$

Finally, we show that the map $\mathcal{K}$ defined by (2.15) and (2.16) has the same equivariance properties with respect to the action of the Weyl group W_G as does the map (2.14). Note that under the identification (2.1), the action of W_G on $H_T^*(M)$ becomes the action

$$w(f_1 \otimes f_2) = f_1 \otimes wf_2 ,$$

since in the right hand side of (2.1), the first factor is $H_G^*(M)$, so W_G acts trivially on it. In particular, the ring of W_G -invariants in $H_T^*(M)$ is

$$\mathbb{S}(\mathfrak{t}^*)^{W_K} \otimes_{\mathbb{S}(\mathfrak{t}^*)^{W_G}} \mathbb{S}(\mathfrak{t}^*)^{W_G} = \mathbb{S}(\mathfrak{t}^*)^{W_K} ,$$

which is consistent with the identifications

$$H_G^*(M) = \mathbb{S}(\mathfrak{t}^*) = \mathbb{S}(\mathfrak{t}^*)^{W_K} = H_T^*(M)^{W_G} . \quad (2.17)$$

On the other hand, the action of W_G on the space

$$H_T^*(M^T) = \text{Maps}(M^T, \mathbb{S}(\mathfrak{t}^*))$$

is just the action

$$(wg)(p) = w(g(w^{-1}p)) ;$$

so to check that the map $\mathcal{K}$ defined by (2.15) and (2.16) is W_G -equivariant, we must show that if

$$g = \mathcal{K}(f_1 \otimes f_2) \quad \text{and} \quad g^w = \mathcal{K}(f_1 \otimes wf_2) ,$$

then for all points $p = \sigma p_0$,

$$g^w(p) = (wg)(p) .$$

However,

$$g^w(p) = (\sigma f_1)(wf_2) = w((w^{-1}\sigma f_1)f_2) = wg(w^{-1}p) = (wg)(p) .$$

Let us now prove that the map $\mathcal{K}$ coincides with the map (2.14). We first note that $\mathcal{K}$ is a morphism of $\mathbb{S}(\mathfrak{t}^*)$ -modules. For $f \in \mathbb{S}(\mathfrak{t}^*)$,

$$\mathcal{K}(f_1 \otimes f_2 f) = \mathcal{K}(f_1 \otimes f_2) f .$$

Thus, it suffices to verify that $\mathcal{K}$ agrees with the map (2.14) on elements of the form $f_1 \otimes 1$. That is, in view of the identification (2.17), it suffices to show that $\mathcal{K}$, restricted to $\mathbb{S}(\mathfrak{t}^*)^{W_K} \otimes 1$, agrees with the map (2.14), restricted to $H_T^*(M)^{W_G}$. However, if $f \in H_T^*(M)^{W_G}$, then $i^*f \in H_T^*(M^T)^{W_G}$, so it

suffices to show that i^*f and $\mathcal{K}(f \otimes 1)$ coincide at p_0 , the identity coset of $M = G/K$. This is equivalent to showing that in the diagram below

$$\begin{array}{ccccc} H_G^*(M) & \longrightarrow & H_K^*(M) & \longrightarrow & H_K^*(p_0) \\ \downarrow & & & & \downarrow \\ \mathbb{S}(\mathfrak{k}^*)^{W_K} & \longrightarrow & & \longrightarrow & \mathbb{S}(\mathfrak{k}^*)^{W_K} \end{array}$$

the bottom arrow is the identity map. However, the bottom arrow is clearly the identity on $\mathbb{S}^0(\mathfrak{k}^*)^K = \mathbb{C}$ and the two maps on the top line are $\mathbb{S}(\mathfrak{k}^*)^K$ -module morphisms. $\square$

3. ALMOST COMPLEX STRUCTURES AND AXIAL FUNCTIONS

3.1. Axial functions. A G -invariant almost structure on $M = G/K$ is determined by an almost complex structure on the tangent space $T_{p_0}M$,

$$J_{p_0} : T_{p_0}M \simeq \mathfrak{g}/\mathfrak{k} \rightarrow \mathfrak{g}/\mathfrak{k} .$$

For an arbitrary point $gp_0 \in M$, the almost complex structure on

$$T_{gp_0}M = (dL_g)_{p_0}(T_{p_0}M)$$

is given by

$$J_{gp_0}((dL_g)_{p_0}(X)) = (dL_g)_{p_0}(J_{p_0}(X)) ,$$

for all $X \in \mathfrak{g}/\mathfrak{k}$. This definition is independent on the representative g chosen if and only if J_{p_0} is K -invariant. Therefore G -invariant almost complex structures on G/K are in one to one correspondence to K -invariant almost complex structures on $\mathfrak{g}/\mathfrak{k}$.

If $M = G/K$ has a G -invariant almost complex structure, then the isotropy representations of T on $T_{p_0}M$ is a complex representation, and therefore its weights are well-defined (not just well-defined up to sign). Let

$$T_{p_0}M = \mathfrak{g}/\mathfrak{k} = \bigoplus_{[\beta]} V_{[\beta]}$$

be the root space decomposition of $\mathfrak{g}/\mathfrak{k}$. Then $V_{[\beta]}$ is a one-dimensional complex representation of T ; let $\tilde{\beta} \in \{\pm\beta\}$ be the weight of this complex representation:

$$\exp t \cdot X_{\tilde{\beta}} = e^{i\tilde{\beta}(t)} X_{\tilde{\beta}} \quad , \text{ for all } t \in \mathfrak{t} .$$

Thus, the map

$$s : \Delta_{G,K}/\pm 1 \rightarrow \Delta_{G,K} , \quad s([\beta]) = \tilde{\beta} , \quad (3.1)$$

is a W_K -equivariant right inverse of the projection $\Delta_{G,K} \rightarrow \Delta_{G,K}/\{\pm 1\}$. Let $\Delta_0 \subset \Delta_{G,K}$ be the image of s .

The existence of a map (3.1) is equivalent to the condition

$$w\alpha \neq -\alpha \quad , \quad \text{for all } w \in W_K , \quad \alpha \in \Delta_{G,K} = \Delta_G - \Delta_K , \quad (3.2)$$

hence (3.2) is a necessary condition for the existence of a G -invariant almost complex structure on M . We will see in the next section that this condition is also sufficient.

We can now define a labelling of the *oriented* edges, E_Γ , of the GKM graph Γ , as follows. Let $[w] \in W_G/W_K$ be a vertex of the graph and let $e = ([w], [w\sigma_\beta])$ be an oriented edge of the graph, with $\beta \in \Delta_0$. This edge corresponds to the subspace $V_{[w\beta]}$ (see (2.10)) in the decomposition

$$T_{[w]}M = \bigoplus_{\beta \in \Delta_0} V_{[w\beta]} ,$$

and the G -invariance of the almost complex structure implies that T acts on $V_{[w\beta]}$ with weight $w\beta$. We define $\alpha : E_\Gamma \rightarrow \mathfrak{t}^*$ by

$$\alpha([w], [w\sigma_\beta]) = w\beta , \text{ for all } \beta \in \Delta_0, w \in W_G . \quad (3.3)$$

Theorem 3.1. *The map $\alpha : E_\Gamma \rightarrow \mathfrak{t}^*$ has the following properties:*

1. *If e_1 and e_2 are two oriented edges with the same initial vertex, then $\alpha(e_1)$ and $\alpha(e_2)$ are linearly independent;*
2. *If e is an oriented edge and $\bar{e}$ is the same edge, with the opposite orientation, then $\alpha(\bar{e}) = -\alpha(e)$;*
3. *If e and e' are oriented edge with the same initial vertex, and if $e'' = \theta_e(e')$, then $\alpha(e'') - \alpha(e')$ is a multiple of $\alpha(e)$.*

Proof. The first assertion is a consequence of the fact that the only multiples of a root α that are roots are $\pm\alpha$.

If e is the oriented edge that joins $[w]$ to $[w\sigma_\beta]$ and that is labelled by $w\beta \in w\Delta_0$, then

$$\alpha(\bar{e}) = (w\sigma_\beta)(\beta) = -w\beta = -\alpha(e) .$$

Finally, if e joins $[w]$ to $[w\sigma_\beta]$ and if e' joins $[w]$ to $[w\sigma_\gamma]$ (with $\beta, \gamma \in \Delta_0$), then e'' joins $[w\sigma_\beta]$ to $[w\sigma_\beta\sigma_\gamma]$, and

$$\alpha(e'') - \alpha(e) = w\sigma_\beta\gamma - w\gamma = w(\sigma_\beta\gamma - \gamma) = -\langle \gamma, \beta \rangle w\beta = -\langle \gamma, \beta \rangle \alpha(e) .$$

□

Equivalently, Theorem 3.1 says that $\alpha : E_\Gamma \rightarrow \mathfrak{t}^*$ is an *axial function* compatible with the connection θ , in the sense of [GZ1].

3.2. Invariant almost complex structures. As we have seen in Section 3.1, (3.2) is a necessary condition for the existence of a G -invariant almost complex structure on $M = G/K$; in this section we show that it is also a sufficient condition.

Theorem 3.2. *If the condition*

$$w\alpha \neq -\alpha \quad , \quad \text{for all } w \in W_K , \alpha \in \Delta_{G,K} = \Delta_G - \Delta_K ,$$

is satisfied, then M admits a G -invariant almost complex structure.

Proof. Consider the complex representation of K on $(\mathfrak{g}/\mathfrak{k})_{\mathbb{C}} = \mathfrak{g}_{\mathbb{C}}/\mathfrak{k}_{\mathbb{C}}$ and let

$$(\mathfrak{g}/\mathfrak{k})_{\mathbb{C}} = \bigoplus_j V_j$$

be the decomposition into irreducible representations; $(\mathfrak{g}/\mathfrak{k})_{\mathbb{C}}$ is self dual, hence

$$\bigoplus_j V_j = (\mathfrak{g}/\mathfrak{k})_{\mathbb{C}} = (\mathfrak{g}/\mathfrak{k})_{\mathbb{C}}^* = \bigoplus_j V_j^* = \bigoplus_j \overline{V_j}$$

Therefore $\overline{V_j} = V_l$ for some l . If α is a highest weight of V_j , then condition (3.2) implies that $-\alpha$ is not a weight of V_j ; however, $-\alpha$ is a weight of $\overline{V_j}$, hence $\overline{V_j} \neq V_j$. Therefore

$$(\mathfrak{g}/\mathfrak{k})_{\mathbb{C}} = \bigoplus_j (V_j \oplus \overline{V_j}) = U \oplus \overline{U}$$

as complex K -representations, and this induces a K -invariant almost complex structure

$$J : \mathfrak{g}/\mathfrak{k} \rightarrow \mathfrak{g}/\mathfrak{k}$$

as follows: If $x \in \mathfrak{g}/\mathfrak{k}$, then there exists a unique $y \in \mathfrak{g}/\mathfrak{k}$ such that $x + iy \in U$, and we define $J(x) = y$. As we have shown before, this is equivalent to the existence of a G -invariant almost complex structure on M . $\square$

An alternative way of proving Theorem 3.2 is to observe that the condition (3.2) is equivalent to the existence of a W_K -equivariant section $s : \Delta_{G,K}/\pm 1 \rightarrow \Delta_{G,K}$. Let s be such a section and let $\Delta_0 \subset \Delta_G - \Delta_K$ be the image of s . Then (see (2.5))

$$\mathfrak{g}/\mathfrak{k} = \bigoplus_{\alpha \in \Delta_0} V_{[\alpha]}$$

and one can define a K -invariant almost complex structure J by requiring that for each $\alpha \in \Delta_0$, J acts on $V_{[\alpha]}$ by

$$J \begin{pmatrix} X_{\alpha} \\ X_{-\alpha} \end{pmatrix} = \begin{pmatrix} X_{-\alpha} \\ -X_{\alpha} \end{pmatrix}. \quad (3.4)$$

4. MORSE THEORY ON THE GKM GRAPH

4.1. Betti numbers. Henceforth we assume that M admits a G -invariant almost complex structure, determined (see (3.4)) by the image $\Delta_0 \subset \Delta_{G,K}$ of a section $s : \Delta_{G,K}/\pm 1 \rightarrow \Delta_{G,K}$. Let Γ be the GKM graph of M and let

$$\alpha : E_{\Gamma} \rightarrow \mathfrak{t}^*$$

be the axial function (3.3). Then the edges whose initial vertex is the identity coset in W_G/W_K are labelled by vectors in Δ_0 .

Let $\xi \in \mathfrak{t}$ be a regular element of $\mathfrak{t}$, *i.e.*

$$\beta(\xi) \neq 0 \quad , \quad \text{for all } \beta \in \Delta_G \subset \mathfrak{t}^* .$$

For a vertex $[w] \in W_G/W_K$, define the *index* of $[w]$ to be

$$\text{ind}_{[w]} = \#\{e \in E_{[w]} ; \alpha(e)(\xi) < 0\} ,$$

and for each $k \geq 0$, let the k -th *Betti number* of Γ be defined by

$$\beta_k(\Gamma) = \#\{[w] \in W_G/W_K ; \text{ind}_{[w]} = k\} .$$

The index of a vertex obviously depends on ξ , but the Betti numbers do not.

Theorem 4.1. [GZ1] *The Betti numbers $\beta_k(\Gamma)$ are combinatorial invariants of Γ (i.e. are independent of ξ).*

In general these Betti numbers are not equal to the Betti numbers

$$\beta_{2k}(M) = \dim H^{2k}(M)$$

of $M = G/K$; however, we show in the next section that there is a large class of homogeneous spaces for which they are equal. One should note that $\beta_{2k}(M)$ is the dimension of the *ordinary* cohomology of M as a *vector space*, while $\beta_k(\Gamma)$ counts the number of generators of degree $2k$ of the *equivariant* cohomology ring of M , as a *free module* over the symmetric algebra $\mathbb{S}(\mathfrak{t}^*)$.

4.2. Morse functions. Let $\xi \in \mathfrak{t}$ be a regular element.

Definition 4.1. *A function $f : W_G/W_K \rightarrow \mathbb{R}$ is called a Morse function compatible with ξ if for every oriented edge $e = ([w], [w'])$ of the GKM graph, the condition $f([w']) > f([w])$ is satisfied whenever $\alpha(e)(\xi) > 0$.*

Morse function do not always exist; however, there is a simple necessary and sufficient condition for the existence of a Morse function: Every regular element $\xi \in \mathfrak{t}$ determines an orientation o_ξ of the edges of Γ : an edge $e \in E_\Gamma$ points upward (with respect to ξ) if $\alpha_e(\xi) > 0$, and points downward if $\alpha_e(\xi) < 0$. The associated directed graph (Γ, o_ξ) is the graph with all upward-pointing edges.

Proposition 4.1. *There exists a Morse function compatible with ξ if and only if the directed graph (Γ, o_ξ) has no cycles.*

4.3. Invariant complex structures. In this section we show that the existence of Morse functions on the GKM graph, which is a combinatorial condition, has geometric implications for the space $M = G/K$.

Theorem 4.2. *The GKM graph (Γ, α) admits a Morse function compatible with a regular $\xi \in \mathfrak{t}$ if and only if the almost complex structure determined by α is a K -invariant complex structure on M . Moreover, if this is the case, then the combinatorial Betti numbers agree with the topological Betti numbers. That is,*

$$b_k(\Gamma) = b_{2k}(M) \quad .$$

Proof. Let $f : W_G/W_K \rightarrow \mathbb{R}$ be a Morse function compatible with ξ , and let $[w]$ be a vertex of the GKM graph where f attains its minimum. If we replace ξ by $w^{-1}(\xi)$ and f by $(w^{-1})^*f$, then the minimum of this new function is p_0 . Thus, without loss of generality, we may assume that the minimum vertex $[w]$ is the identity coset in W_G/W_K . Then

$$\Delta_0 = \{\beta \in \Delta_{G,K} ; \beta(\xi) > 0\} ,$$

hence Δ_0 is the intersection of $\Delta_{G,K}$ with the positive Weyl chamber determined by ξ . Let

$$\mathfrak{p} = \mathfrak{k}_{\mathbb{C}} \oplus \left(\bigoplus_{\beta \in \Delta_0} \mathfrak{g}_{\beta} \right) .$$

Then $\mathfrak{p}$ is a parabolic subalgebra of $\mathfrak{g}_{\mathbb{C}}$, hence the almost complex structure determined by α is actually a complex structure.

If $G_{\mathbb{C}}$ is the simply connected Lie group with Lie algebra $\mathfrak{g}_{\mathbb{C}}$ and if P is the Lie subgroup of $G_{\mathbb{C}}$ corresponding to $\mathfrak{p}$, then

$$M = G/K = G_{\mathbb{C}}/P ,$$

hence M is a coadjoint orbit of G . Then M is a Hamiltonian T -space and the GKM graph of M is the 1-skeleton of the moment polytope, and therefore the combinatorial Betti numbers agree with the topological Betti numbers.

On the other hand, if the almost complex structure is integrable then $\mathfrak{p}$ is a parabolic subalgebra of $\mathfrak{g}_{\mathbb{C}}$, $M = G/K \subset \mathfrak{g}^*$ is a coadjoint orbit of G , and for a generic direction $\xi \in \mathfrak{t} \subset \mathfrak{g}$, the map $f : W_G/W_K \rightarrow \mathbb{R}$ given by

$$f([w]) = \langle [w], \xi \rangle$$

(with $W_G/W_K \rightarrow G/K \rightarrow \mathfrak{g}^*$) is a Morse function on the GKM graph compatible with ξ . $\square$

5. EXAMPLES

5.1. Non-existence of almost complex structures. Let G be a compact Lie group such that $\mathfrak{g}_{\mathbb{C}}$ is the simple Lie algebra of type B_2 . Let $\alpha_1, \alpha_1 + \alpha_2$




FIGURE 1.

be the short positive roots and let $\alpha_2, \alpha_2 + 2\alpha_1$ be the long positive roots.

Let K be the subgroup of G corresponding to the root system consisting of the short roots. Then $\mathfrak{k}_{\mathbb{C}} = D_2 = A_1 \times A_1$ and $K \simeq SU(2) \times SU(2)$. The quotient W_G/W_K has two classes: the class of $\sigma_{\alpha_1} \in W_K$ and the class of $\sigma_{\alpha_2} \in W_G - W_K$.

The GKM graph Γ has two vertices, joined by two edges, and the edges are labelled by $[\alpha_2], [\alpha_2 + 2\alpha_1] \in \Delta_{G,K}/\pm 1$. If $w = \sigma_{\alpha_1 + \alpha_2} \sigma_{\alpha_1} \in W_K$, then $w\alpha_2 = -\alpha_2$ and $\alpha_2 \in \Delta_{G,K}$, hence one can't define an axial function on Γ . In this example, $G/K = S^4$, which does not admit an almost complex structure.

5.2. Non-existence of Morse functions. Let G be a compact Lie group such that $\mathfrak{g}_{\mathbb{C}}$ is the simple Lie algebra of type G_2 . Let $\alpha_1, \alpha_1 + \alpha_2$, and $2\alpha_1 + \alpha_2$ be the short positive roots and let $\alpha_2, 2\alpha_2 + 3\alpha_1, \alpha_2 + 3\alpha_1$ be the long positive roots. Let K be the subgroup of G corresponding to the root system consisting of the short roots. Then $\mathfrak{k}_{\mathbb{C}} = A_2$ and $K \simeq SU(3)$. The quotient W_G/W_K has two classes: the class of $\sigma_{\alpha_1} \in W_K$ and the class of $\sigma_{\alpha_2} \in W_G - W_K$.




FIGURE 2.

The GKM graph Γ has two vertices, joined by three edges, and the edges are labelled by $[\alpha_2], [2\alpha_2 + 3\alpha_1], [\alpha_2 + 3\alpha_1] \in \Delta_{G,K}/\pm 1$. There are two W_K -equivariant sections of the projection $\Delta_{G,K} \rightarrow \Delta_{G,K}/\pm 1$, corresponding to $\{\alpha_2, \alpha_2 + 3\alpha_1, -2\alpha_2 - 3\alpha_1\}$ and $\{-\alpha_2, -\alpha_2 - 3\alpha_1, 2\alpha_2 + 3\alpha_1\}$. If

$$\Delta_0 = \{\alpha_2, \alpha_2 + 3\alpha_1, -2\alpha_2 - 3\alpha_1\},$$

then the axial function is shown in Figure 2 and there is no Morse function on Γ : the corresponding almost complex structure is not integrable. In this example, $G/K = S^6$, which admits an almost complex structure, but no invariant complex structure.

5.3. The existence of several almost complex structures. Let $G = SU(3)$ and $K = T$. Then the homogeneous space G/K is the manifold of complete flags in $\mathbb{C}^3$. The root system of G is A_2 , with positive roots α_1, α_2 ,

and $\alpha_1 + \alpha_2$ of equal length. The Weyl group of G is $W_G = S_3$, the group of permutations of $\{1, 2, 3\}$, and $W_K = 1$, hence $W_G/W_K = W_G = S_3$.

The GKM graph is the bi-partite graph $K_{3,3}$: it has 6 vertices and each vertex has 3 edges incident to it, labelled by $[\alpha_1]$, $[\alpha_2]$, and $[\alpha_1 + \alpha_2]$. There are 2^3 possible W_K -invariant sections, hence eight G -invariant almost complex structures on G/K . If

$$\Delta_0 = \{\alpha_1, \alpha_2, \alpha_1 + \alpha_2\} ,$$

then the corresponding almost complex structure is integrable and there is a Morse function on Γ compatible with $\xi \in \mathfrak{t}$ such that both $\alpha_1(\xi)$, and $\alpha_2(\xi)$ are positive. This Morse function is given by $f(w) = \ell(w)$ where $\ell(w)$ is the length of w , *i.e.*, in this case, the number of inversions in w .




FIGURE 3.

However, if

$$\Delta_0 = \{\alpha_1, \alpha_2, -\alpha_1 - \alpha_2\} ,$$

then the corresponding almost complex structure is not integrable and there is no Morse function on (Γ, α) : for every vertex w of Γ , there exist three edges e_1, e_2 , and e_3 , going out of w , such that

$$\alpha_{e_1} + \alpha_{e_2} + \alpha_{e_3} = 0 ,$$

hence there is no vertex of Γ on which a Morse function compatible with some $\xi \in \mathfrak{t}$ can achieve its minimum.

REFERENCES

- [Bo] A. Borel, *Seminar on transformation groups*, Ann. of Math. Stud. **46**, Princeton Univ. Press, Princeton, NJ 1960.
- [FH] W. Fulton and J. Harris, *Representation Theory*, Springer, New York 1991.
- [GHV2] W. Greub, S. Halperin, and R. Vanstone, *Connections, Curvature, and Cohomology*, vol. II, Academic Press, 1973.

- [GHV3] W. Greub, S. Halperin, and R. Vanstone, *Connections, Curvature, and Cohomology*, vol. III, Academic Press, 1976.
- [GKM] M. Goresky, R. Kottwitz and R. MacPherson, *Equivariant cohomology, Koszul duality, and the localization theorem*, Invent. Math. **131** (1998), no. 1, pp. 25-83.
- [Go] R. Goldin, *The Cohomology of Weight Varieties*, Ph.D. thesis, MIT 1999.
- [GS] V. Guillemin and S. Sternberg, *Supersymmetry and equivariant de Rham cohomology*, Springer Verlag, Berlin 1999.
- [GZ1] V. Guillemin and C. Zara, *One-skeleta, Betti numbers and Equivariant Cohomology*, Duke Math. J., **107** (2001), no. 2, pp. 283-349.
- [Hu] J. Humphreys, *Reflection groups and Coxeter groups*, Cambridge Univ. Press, 1990.
- [Ho] T. Holm, *Homogeneous spaces, equivariant cohomology, and graphs*, Ph.D. thesis, MIT 2002.
- [V] D. Vogan, personal communication.

DEPARTMENT OF MATHEMATICS, MIT, CAMBRIDGE, MA 02139
E-mail address: `vwg@math.mit.edu`

DEPARTMENT OF MATHEMATICS, MIT, CAMBRIDGE, MA 02139
E-mail address: `tsh@math.mit.edu`

DEPARTMENT OF MATHEMATICS, YALE UNIVERSITY, NEW HAVEN, CT 06520
E-mail address: `zara@math.yale.edu`